



## Main sol. J950

Na <sub>2</sub> SO <sub>4</sub>	1.00	g
NaCl	30.00	g
KCl	0.70	g
MgCl <sub>2</sub> x 6 H <sub>2</sub> O	5.00	g
CaCl <sub>2</sub> x 2 H <sub>2</sub> O	1.00	g
K <sub>2</sub> HPO <sub>4</sub>	0.40	g
KH <sub>2</sub> PO <sub>4</sub>	0.40	g
NH <sub>4</sub> Cl	0.50	g
Fe(NH <sub>4</sub> ) <sub>2</sub> (SO <sub>4</sub> ) <sub>2</sub> x 6 H <sub>2</sub> O	2.00	mg
Yeast extract (BD-Difco)	3.00	g
Tryptone (BD-Difco)	4.00	g
Sodium acetate	6.80	g
L-Cysteine HCl x H <sub>2</sub> O	0.60	g
<b>Trace element solution SL-10</b>	1.00	ml
Resazurin	1.00	mg
Distilled water	960.00	ml

Mix components thoroughly and adjust pH to 6.8. Distribute the medium into culture vessels under a N<sub>2</sub>-CO<sub>2</sub> (9:1, v/v) gas mixture, seal with butyl rubber stoppers and autoclave. After cooling, add the following solution (per 960 ml) from anaerobic stock (filter-sterilized)

NaHCO <sub>3</sub> (5%)	40.00	ml
-------------------------	-------	----