

104: PYG MEDIUM (modified)

This recipe contains strain-specific modifications for *Coprobacillus tertius* DSM 114538 *

Final pH: 7.2

Final volume: 1000 ml

Trypticase peptone	5.00	g
Peptone	5.00	g
Yeast extract	10.00	g
Beef extract	5.00	g
Glucose	5.00	g
K ₂ HPO ₄	2.00	g
Tween 80	1.00	ml
Salt solution	40.00	ml
Resazurin	1.00	mg
Vitamin K₁ solution	0.20	ml
Haemin solution	10.00	ml
L-Cysteine HCl x H ₂ O	0.50	g
Distilled water	950.00	ml

The vitamin K₁, haemin solution and the cysteine are added after the medium has been boiled and cooled under CO₂. Adjust pH to 7.2 using 8 N NaOH. Distribute under N₂ and autoclave.

* dilute the medium 1:10

Salt solution (from medium 104)

CaCl ₂ x 2 H ₂ O	0.25	g
MgSO ₄ x 7 H ₂ O	0.50	g
K ₂ HPO ₄	1.00	g
KH ₂ PO ₄	1.00	g
NaHCO ₃	10.00	g
NaCl	2.00	g
Distilled water	1000.00	ml

Haemin solution (from medium 78)

Haemin	50.00	mg
NaOH (1 N)	1.00	ml
Distilled water	100.00	ml

Dissolve 50 mg haemin in 1 ml 1 N NaOH; make up to 100 ml with distilled water and filter sterilize. Store refrigerated.

104: PYG MEDIUM (modified)

Vitamin K₁ solution (from medium 78)

Vitamin K ₁	0.10	ml
Ethanol (95 %)	20.00	ml

Dissolve 0.1 ml of vitamin K₁ in 20 ml 95% ethanol and filter sterilize. Store refrigerated in a brown bottle.