## **Microorganisms**



## **104: PYG MEDIUM (modified)**

This recipe contains strain-specific modifications for Veillonella montpellierensis DSM 17217 \*

Final pH: 7.2 Final volume: 1000 ml

Trypticase peptone	5.00	g
Peptone	5.00	g
Yeast extract	10.00	g
Beef extract	5.00	g
Glucose	5.00	g
K <sub>2</sub> HPO <sub>4</sub>	2.00	g
Tween 80	1.00	ml
Salt solution	40.00	ml
Resazurin	1.00	mg
Vitamin K <sub>1</sub> solution	0.20	ml
Haemin solution	10.00	ml
L-Cysteine HCl x $H_2O$	0.50	g
Putrescine	10.00	µl/ml
Sodium lactate	10.00	µl/ml
Distilled water	950.00	ml

The vitamin  $K_1$ , haemin solution and the cysteine are added after the medium has been boiled and cooled under  $CO_2$ . Adjust pH to 7.2 using 8 N NaOH. Distribute under  $N_2$  and autoclave.

## \* Plus 10µl/ml putrescine (0.03%) plus 10 µl/ml sodium lactate (50%)

Salt solution (from medium 104)		
$CaCl_2 \times 2 H_2O$	0.25	g
$MgSO_4 \times 7 H_2O$	0.50	g
K <sub>2</sub> HPO <sub>4</sub>	1.00	g
KH <sub>2</sub> PO <sub>4</sub>	1.00	g
NaHCO <sub>3</sub>	10.00	g
NaCl	2.00	g
Distilled water	1000.00	ml
Hoomin colution (from modium 79)		
Haemin solution (from medium 78) Haemin	50.00	ma
		mg
NaOH (1 N)	1.00	ml
Distilled water	100.00	ml

## Microorganisms

104: PYG MEDIUM (modified)



Dissolve 50 mg haemin in 1 ml 1 N NaOH; make up to 100 ml with distilled water and filter sterilize. Store refrigerated.

Vitamin K <sub>1</sub> solution (from medium 78)		
Vitamin K <sub>1</sub>	0.10	ml
Ethanol (95 %)	20.00	ml

Dissolve 0.1 ml of vitamin  $K_1$  in 20 ml 95% ethanol and filter sterilize. Store refrigerated in a brown bottle.