

135a: ACETOBACTERIUM MEDIUM (MARINE)

This recipe contains strain-specific modifications for *Salileptolyngbya* sp. DSM 2396 *

Final pH: 7.0 - 7.2

Final volume: 1021 ml

NaCl	20.00	g
NH ₄ Cl	1.00	g
KH ₂ PO ₄	0.33	g
K ₂ HPO ₄	0.45	g
MgSO ₄ x 7 H ₂ O	0.10	g
Modified Wolin's mineral solution	20.00	ml
Yeast extract	2.00	g
Resazurin (0.1% w/v)	0.50	ml
NaHCO ₃	5.00	g
Ethylene glycol	1.25	g
Wolin's vitamin solution (10x)	1.00	ml
L-Cysteine HCl x H ₂ O	0.50	g
Na ₂ S x 9 H ₂ O	0.50	g
Distilled water	1000.00	ml

Dissolve ingredients except bicarbonate, ethylene glycol, vitamins, cysteine and sulfide, bring to the boil and cool to room temperature under 80% N₂ and 20% CO₂ gas mixture. Add bicarbonate (solid) and equilibrate the medium with the gas until a pH of around 7.4 is reached. Then distribute under the same gas atmosphere into anoxic Hungate-type tubes or serum vials and autoclave. Before use add ethylene glycol, vitamins (sterilized by filtration), cysteine and sulfide from sterile anoxic stock solutions prepared under 100% N₂ gas. Adjust pH of complete medium to 7.0 - 7.2, if necessary.

*

Modified Wolin's mineral solution (from medium 141)

Nitrolotriacetic acid	1.50	g
MgSO ₄ x 7 H ₂ O	3.00	g
MnSO ₄ x H ₂ O	0.50	g
NaCl	1.00	g
FeSO ₄ x 7 H ₂ O	0.10	g
CoSO ₄ x 7 H ₂ O	0.18	g
CaCl ₂ x 2 H ₂ O	0.10	g
ZnSO ₄ x 7 H ₂ O	0.18	g
CuSO ₄ x 5 H ₂ O	0.01	g
AlK(SO ₄) ₂ x 12 H ₂ O	0.02	g
H ₃ BO ₃	0.01	g

Na ₂ MoO ₄ x 2 H ₂ O	0.01	g
NiCl ₂ x 6 H ₂ O	0.03	g
Na ₂ SeO ₃ x 5 H ₂ O	0.30	mg
Na ₂ WO ₄ x 2 H ₂ O	0.40	mg
Distilled water	1000.00	ml

First dissolve nitrilotriacetic acid and adjust pH to 6.5 with KOH, then add minerals. Adjust final to pH 7.0 with KOH.

Wolin's vitamin solution (10x) (from medium 120)

Biotin	20.00	mg
Folic acid	20.00	mg
Pyridoxine hydrochloride	100.00	mg
Thiamine HCl	50.00	mg
Riboflavin	50.00	mg
Nicotinic acid	50.00	mg
Calcium D-(+)-pantothenate	50.00	mg
Vitamin B ₁₂	1.00	mg
p-Aminobenzoic acid	50.00	mg
(DL)-alpha-Lipoic acid	50.00	mg
Distilled water	1000.00	ml