

141c: METHANOCOCCOIDES MEDIUM

This recipe contains strain-specific modifications for *Methanosarcina mazei* DSM 3318 *

Final pH: * 6.8 - 7.2

Final volume: 1013 ml

KCl	0.34	g
MgCl ₂ x 6 H ₂ O	4.00	g
MgSO ₄ x 7 H ₂ O	3.45	g
NH ₄ Cl	0.25	g
CaCl ₂ x 2 H ₂ O	0.14	g
K ₂ HPO ₄	0.14	g
NaCl	18.00	g
Modified Wolin's mineral solution	10.00	ml
Fe(NH ₄) ₂ (SO ₄) ₂ x 6 H ₂ O (0.1% w/v)	2.00	ml
Na-acetate	1.00	g
Yeast extract (OXOID)	2.00	g
Trypticase peptone (BD BBL)	2.00	g
Sodium resazurin (0.1% w/v)	0.50	ml
NaHCO ₃	5.00	g
Trimethylammonium chloride	5.00	g
Wolin's vitamin solution (10x)	1.00	ml
L-Cysteine HCl x H ₂ O	0.50	g
Na ₂ S x 9 H ₂ O	0.50	g
Methanol	5.00	ml
Distilled water	1000.00	ml

Dissolve ingredients (except bicarbonate, trimethylammonium chloride, vitamins, cysteine and sulfide), sparge medium with 80% N₂ and 20% CO₂ gas mixture for 30 - 45 min to make it anoxic. Add and dissolve bicarbonate and adjust pH to 6.8, then dispense medium under 80% N₂ and 20% CO₂ gas atmosphere into anoxic Hungate-type tubes or serum vials to 30% of their volume and autoclave. After sterilization add trimethylammonium chloride, cysteine and sulfide from sterile anoxic stock solutions autoclaved under 100% N₂ gas atmosphere. Vitamins are prepared under 100% N₂ gas atmosphere and sterilized by filtration. Adjust pH of complete medium to 7.0 - 7.4, if necessary.

* Replace trimethylammonium chloride with 10.00 ml/l methanol solution (50% v/v) added to the autoclaved medium from a sterile anoxic stock solution. Adjust pH of complete medium to 6.8 - 7.2.

Modified Wolin's mineral solution (from medium 141)

Nitrilotriacetic acid	1.50	g
MgSO ₄ x 7 H ₂ O	3.00	g

141c: METHANOCOCCOIDES MEDIUM

MnSO ₄ x H ₂ O	0.50	g
NaCl	1.00	g
FeSO ₄ x 7 H ₂ O	0.10	g
CoSO ₄ x 7 H ₂ O	0.18	g
CaCl ₂ x 2 H ₂ O	0.10	g
ZnSO ₄ x 7 H ₂ O	0.18	g
CuSO ₄ x 5 H ₂ O	0.01	g
AlK(SO ₄) ₂ x 12 H ₂ O	0.02	g
H ₃ BO ₃	0.01	g
Na ₂ MoO ₄ x 2 H ₂ O	0.01	g
NiCl ₂ x 6 H ₂ O	0.03	g
Na ₂ SeO ₃ x 5 H ₂ O	0.30	mg
Na ₂ WO ₄ x 2 H ₂ O	0.40	mg
Distilled water	1000.00	ml

First dissolve nitrilotriacetic acid and adjust pH to 6.5 with KOH, then add minerals. Adjust final to pH 7.0 with KOH.

Wolin's vitamin solution (10x) (from medium 120)

Biotin	20.00	mg
Folic acid	20.00	mg
Pyridoxine hydrochloride	100.00	mg
Thiamine HCl	50.00	mg
Riboflavin	50.00	mg
Nicotinic acid	50.00	mg
Calcium D-(+)-pantothenate	50.00	mg
Vitamin B ₁₂	1.00	mg
p-Aminobenzoic acid	50.00	mg
(DL)-alpha-Lipoic acid	50.00	mg
Distilled water	1000.00	ml